

INFANT CASES

| | |
|-----------------------|--|
| Cefotaxime | Dose: 50mg/kg/dose q8 hourly IV |
| Ampicillin | Dose: 150 mg/kg/day divided q6 hourly IV |
| RSI Meds | |
| Lidocaine | Dose: 7.5 - 10.8 mg (1.5mg/kg) IV |
| Atropine | Dose: 0.1-0.5mg 0.02 mg/kg IV |
| Fentanyl | Dose: 14.4 - 21.6 mcg (3 mcg/kg) IV |
| Propofol | Dose: 14.4 - 21.6 mg (3mg/kg) IV |
| Etomidate | Dose: 1.4 - 2.1 mg (0.3mg/kg) IV |
| Ketamine | Dose: 9.6 - 14.4 mg (2mg/kg) IV |
| Succinylcholine | Dose: 7.5 - 10.8 mg (1.5mg/kg) IV |
| Vecuronium | Dose: 0.5 - 0.7 mg (0.1mg/kg) IV |
| Rocuronium | Dose: 4.8 - 7.2 mg (1mg/kg) IV |
| Seizure Meds | |
| Ativan(Lorazepam) | Dose: 0.48 - 0.72 mg (0.1 mg/kg) |
| Diastat | Dose: 0.24 - 0.36mg (0.5mg/kg) rectally |
| Dilantin(Phenytoin) | Dose: 75 - 108 mg (15mg/kg)IV |
| Keppra(Levetiracetam) | Dose: 50mg/kg/IV |
| Valium(Diazepam) | Dose: 0.96 - 1.44 mg (0.2mg/kg) IV |
| Versed(Midazolam) | Dose: 0.5 - 0.7 mg (0.1 mg/kg) IV |
| Phenobarbital | Dose: 96 - 144 mg (20mg/kg) IV |
| Fosphenytoin | Dose: 75 - 108 mg (15mg/kg)IV |

CHILD CASES- CPR

| | |
|--------------------------------|--|
| Amiodarone | Dose: 80 - 120 mg (5 mg/kg) IV |
| Morphine Sulfate | Dose: 1.6 - 2.4 mg (0.1 mg/kg) IV |
| 10% Calcium Chloride | Dose: 3.2 - 4.8 ml (0.2 mL/kg) IV Dose: 320 - 480 mg (20 mg/kg) IV |
| Sodium Bicarbonate 8.4% | Dose: 16 - 24 ml (1 mL/kg) IV Dose: 16 - 24 mEq (1 mEq/kg) IV |
| Ativan(Lorazepam) | Dose: 1.6 - 2.4 mg (0.1 mg/kg) IV |
| Valium(Diazepam) | Dose: 3.2 - 4.8 mg (0.2mg/kg) IV |
| Versed(Midazolam) | Dose: 1.6 - 2.4 mg (0.1 mg/kg) IV |
| Lidocaine 2% | Dose: 16 - 24 mg (1 mg/kg) IV/ETT |
| RSI (premed) Atropine | Dose: 0.32 - 0.48 (0.02 mg/kg) IV/ETT |
| RSI (premed) Fentanyl | Dose: 16 - 24 mcg (1 mcg/kg) IV |
| RSI (sedative)Propofol | Dose: 48 - 72 (3mg/kg) IV |
| RSI (sedative) Etomidate | Dose: 4.8 - 7.2 mg (0.3mg/kg) IV |
| RSI (sedative) Ketamine | Dose: 32 - 48 mg (2mg/kg) IV |
| RSI (paralytic) Succinylcholin | Dose: 24 - 36 (1.5mg/kg) IV |
| RSI (paralytic) Vecuronium | Dose: 1.6 - 2.4 mg (0.1mg/kg) IV |
| RSI (paralytic) Rocuronium | Dose: 16 - 24 mg (1mg/kg) IV |